

**Supplementary Table 1. Baseline Disease Characteristics (Safety Population)**

	MTX				Overall
	10 mg	15 mg	20 mg	25 mg	
	(n=13)	(n=12)	(n=12)	(n=12)	
ACR RA classification, n (%)					
Stage I	2 (15.4)	0 (0.0)	0 (0.0)	1 (8.3)	3 (6.1)
Stage II	5 (38.5)	6 (50.0)	11 (91.7)	8 (66.7)	30 (61.2)
Stage III	6 (46.2)	5 (41.7)	1 (8.3)	2 (16.7)	14 (28.6)
Stage IV	0 (0.0)	1 (8.3)	0 (0.0)	1 (8.3)	2 (4.1)
Functional status, n (%)					
Class I	1 (7.7)	2 (16.7)	2 (16.7)	2 (16.7)	7 (14.3)
Class II	7 (53.8)	9 (75.0)	9 (75.0)	8 (66.7)	33 (67.3)
Class III	5 (38.5)	1 (8.3)	1 (8.3)	2 (16.7)	9 (18.4)
Class IV	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)

ACR, American College of Rheumatology; MTX, methotrexate; RA, rheumatoid arthritis.

**Supplementary Table 2. Adverse Reactions to Methotrexate Treatment (Safety Population; Each Patient Received 1 oral MTX administration, 1 MTXAI delivered to the abdomen, and 1 MTXAI delivered to the thigh)**

	N*	Comments
<b>Adverse events</b>		
SAEs	2	<ul style="list-style-type: none"> <li>• 1 fatal MI (study day 6): 79-year-old man with prior cardiac history (MTXAI 25 mg abdomen)</li> <li>• Sick sinus syndrome (after trial concluded): 72-year-old man with history of CAD (MTXAI 15 mg thigh)</li> </ul>
At least 1 TEAE <sup>†</sup>	3	Fatigue, nausea, worsening of RA <sup>‡</sup>
<b>Injection site reactions</b>		
Erythema	2	Grade 1

\*Number of patients.

<sup>†</sup>Excluding TEAEs classified as SAEs.

<sup>‡</sup>Also led to discontinuation.

CAD, coronary artery disease; MI, myocardial infarction; MTXAI, methotrexate auto-injector; SAE, serious adverse event; TEAE, treatment-emergent adverse event.